

Modelling of the Contribution of Different Blended Fertilizers to Potato Production in Basketo Special Woreda, Gamo-gofa Zone, Ethiopia

Metadel Minaye Getahun¹, Zeleke Eromo Ewunetu², Kindu Kebede Gebre^{*3}, Zelalem Abera Amare⁴, Tarekegn Keriso Menisho⁵

¹Arba Minch Agricultural Research Center, Southern Agricultural Research Institute, Hawassa, Ethiopia.
Email: siketmeta21@gmail.com | ORCID: <https://orcid.org/0000-0001-7209-4542>

²Arba Minch Agricultural Research Center, Southern Agricultural Research Institute, Hawassa, Ethiopia.
Email: dreamyzed2005@gmail.com | ORCID: <https://orcid.org/0000-0002-5452-3179>

³Department of Statistics, College of Computing and Informatics, Haramaya University, Dire Dawa, Ethiopia.
Email: m7.kebede@gmail.com | ORCID: <https://orcid.org/0000-0002-5337-4343>

⁴University Laboratory Management Directorate Office, Central Laboratory, Haramaya University, Dire Dawa, Ethiopia. Email: aberazelalem2@gmail.com | ORCID: <https://orcid.org/0000-0002-6017-2352>

⁵Arba Minch Agricultural Research Center, Southern Agricultural Research Institute, Hawassa, Ethiopia.
Email: yedideya04@gmail.com | ORCID: <https://orcid.org/0000-0002-7099-2020>

*Corresponding Author

How to cite this paper: Getahun, M.M., Ewunetu, Z.E., Gebre, K.K., Amare, Z.A. and Menisho, T.K. (2022). Modelling of the Contribution of Different Blended Fertilizers to Potato Production in Basketo Special Woreda, Gamo-gofa Zone, Ethiopia. *Grassroots Journal of Natural Resources*, 5(2): 1-11. Doi: <https://doi.org/10.33002/nr2581.6853.050201>

Received: 11 January 2022

Reviewed: 16 February 2022

Provisionally Accepted: 29 March 2022

Revised: 30 April 2022

Finally Accepted: 17 May 2022

Published: 27 June 2022

Copyright © 2022 by author(s)

This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).
<http://creativecommons.org/licenses/by/4.0/>



Open Access

Editor-in-Chief
Prof. Dr. G. Poyyamoli
Executive Editor
Dr. Hasrat Arjjumend
Associate Editors
Dr. Maja Manojlovic
Ms. Areej Sabir
Dr. Usongo Patience



Abstract

Potato is important for food security in developing countries such as Ethiopia. The purpose of this study is to evaluate the contribution of varied fertilizers to potato yields at Basketo Special Woreda, Gamogofa Zone, Ethiopia. To attain this objective, a generalized linear mixed model was used to differentiate the factors that affect the marketable yield of potatoes by considering variability within three replications. In addition, this study used the Kolmogorov-Smirnov test to check the goodness of fit tests of normal distribution. The collected data were analyzed using statistical software, SAS version 9.4. The mean of plant tallness, number of plants per plot, number of plants per slope, and number of tubers per slope were 73.65 (SD = 12.2), 155.2 (SD = 112.2), 3.23 (SD = 1.31), and 10.91 (SD = 3.15.2), respectively. The result reveals that 6% of variation existed in potato yield between plots, and the remaining 94% existed within plots. Apart from a linear mixed model, a random intercept model was applied to this data. The findings reveal that fertilizer types, number of plants per plot, and numbers of tubers per hill were significant at 5% level of significant. Thus, the expected marketable yield of potatoes decreased by 0.7819 tons/hectare when the number of tubers per hill increased by 1 unit. On the other hand, the expected marketable yield of potatoes has increased by 0.0175 tons/hectare when the number of plants per plot increased by 1 unit.

Keywords

Marketable yield; Potato; Fertilizer types; Random Intercept Model; Ethiopia

Introduction

The potato (*Solanum tuberosum*) is an Andean staple meal crop (Rodríguez *et al.*, 2010). Potato is basically produced in Europe and North America. It is the most important meal crop and is used for food intake (Sonnewald *et al.*, 2015). It has significant value in terms of food security in developing countries like Ethiopia (Girma *et al.*, 2020). For instance, farmers in Ethiopia produced potatoes from 0.5 million tons in 2006 to 3.6 million tons in 2016 (Voorra, Larrea and Bermudez, 2020). It has widely been grown in the highlands of Ethiopia. However, the yield of potatoes is low due to improperly used fertilizers (Alemayehu and Jemberie, 2018).

Potato production requires fertilizers like phosphorus (P). Though phosphorus (P), an important nutrient for potato production, is required in excess quantities, potatoes make inefficient use of soil phosphorus (Rosen *et al.*, 2014). In addition, farmers used nitrogen fertilizer to maximize tuber size (Rens *et al.*, 2018). Sulphur fertilizer is also important for increasing shoot and root interiors (Hawkesford and De Kok, 2006). Additionally, sulphur absorption is used for coordination with carbon and nitrogen pathways (Hawkesford and De Kok, 2006).

Many research findings on potato production in Ethiopia have focused on the impact of integrated application of organic and blended mineral fertilizers on potato productivity. For example, a study revealed the impact of natural and inorganic fertilizer application at the potato surrender (Girma *et al.*, 2020). Similarly, an investigation by Girma, Beyene and Biazin (2017, pp.1 & 6) confirmed the influence of phosphorus and potassium fertilizers on potato growth and yield. Another study claimed the reaction of potato to nitrogen and phosphorus (Setu, Dechassa and Alemayehu, 2018). Moreover, another study disclosed the impact of fertilization administration at the trim execution and chemical composition of potato cultivated in Greece (Sebnie, Esubalew and Mengesha, 2021). Most studies have been done on to improve potato production. However, none of them show the effect of fertilizer types within and between plots. Therefore, this study is conducted in assessing the contribution of different blended fertilizers to potato production as fertilizers can have an impact on the environment and production (Briefs, 2019). In doing so, the study used conditional residual plots and Kolmogorov–Smirnov test to check the normality of the error term and goodness of fit, respectively (Massey Jr, 1951; Wang *et al.*, 2018). In addition, in this study, generalized linear mixed models were used to assess the contribution of different blended fertilizer types to potato production and estimate the variation of potato yield within and between plots in the Basketo Special Woreda, Gamogofa Zone, Ethiopia.

Study Design and Methods

Study Area

The field experiment was carried out at Arba Minch Agricultural Research Center, in the Southern Regional State of Ethiopia, during the 2018 and 2019 cropping seasons. Basketo Special Woreda is located in the Gamo-gofa Zone, Ethiopia. It lies at the latitude of 6.25° and a longitude of 36.58333°, with GPS coordinates of 6° 15' North and 36° 35' East.

Experiment Instruments

The potato variety named as "Belete" was planted and harvested by Holeta Rural Investigators and was utilized for the experiment (Tessema, Mohammed and Abebe, 2020). Fertilizers were applied on the sides of potato tubers on the edges. Phosphorus, sulphur, potassium, and boron nutrients were applied at the planting time; nitrogen was applied in split halves during planting and after planting. Potassium chloride (KCl), triple super phosphate (TSP), nitrogen, phosphorus, sulphur and boron (NPSB), and nitrogen, phosphorus, and sulphur (NPS) fertilizers were the nutrient sources.

The field test was laid out at random and consisted of five chemicals with three replicates each. The trial was conducted during the autumn season in 2018 and 2019. Potato tubers were planted in a 30 cm gap with 75 cm row spacing. This study used 15 total plots with a 3.9 m by 3.75 m unit size of the plot area, and each accommodated 5 rows and 13 columns. The distance between and inside a plot was 1 m and 1.5 m, respectively.

The experimental crop variety was an improved variety (Belete). Soil samples (0–20 cm depth) before and after sowing were taken randomly in a zig zag pattern from 10 soil cores at the same depth of 0–20 cm. The soil samples were combined to form a composite 1 kg soil sample in a sterilized soil bag. Soil drying was accomplished by spreading each sample on neat paper to air dry at room temperature, then bagging and freezing for shipping. Analyses of pH, OC TN, P, K, S, and texture parameters of soil samples were analyzed within two weeks. The sample field was laid out in a randomized complete block design having five treatments: NP (165:195), NPS (120:92:17), NPSZnB (120:92:20.06:5.89:0.66), NPKSZnB (120:92:52.17:19.74:6.06:0.88), and control (no fertilizer used) based on agricultural transformation agency's recommendation (Bellete, 2015). The agricultural transformation agency recommends that farmers apply a high amount of NP as compared to other treatments in Baskato Woreda. The first and second-year experiments were undertaken in the main rainy season (Belg from March to May) in 2018 and 2019.

Experiment Procedures and Data Collection

This study is based on an imminent cohort study design using field tests. It is important to connect the sides of potato tubers and secure them with soil to maintain a strategic distance. The suggested rates of P, S, K, Zn and B supplements are used as the chemicals to connect planting, while N is connected in part at half planting and the remaining half after planting was used for almost 21 days. Moreover, in the area, NPSB, NPS, and KCl fertilizers were utilized as supplements (Bellete, 2015).

Surface soil test samples (8 cores at 0–20 cm depth) were taken arbitrarily from the field before the start of the field experiment. The soil tests were composited and arranged for examinations of the soil's physical-chemical properties. Ph (6.08), OC (2.6%), OM (4.6%), TN (0.23%), available P (6.5 mg/l), and soil textural classes were clay loam for the selected farm fields at the soil laboratory of Arba Minch Agricultural Research Center, and Southern Agricultural Institute. A simple random sampling technique was used for plant selection having five plants. The authors counted the whole number of plants per plot, plants per hill, and tubers per hill, and measured the plant height of selected plants. Besides, this study estimated the marketable yield per plot and recorded it by excluding the border row.

Quality of Data Collection

The quality of the data was approved by information controllers who were assigned from the Arba Minch Agricultural Research Center. The information extraction apparatuses and the completeness of the information were controlled by the Arba Minch Agricultural Research Centre. All necessary amendments were made to the ultimate information collection sheet.

Study Variables

The indicator factors were fertilizer sort, number of plants per plot, number of tubers per slope, and plant stature. In addition, the marketable yield of potatoes was the outcome variable of the study.

Statistical Methods of Analysis for Repeated Data

This study used generalized linear mixed models, which are a natural outgrowth of both linear mixed models and generalized linear models. The generalized linear mixed model can be measured repeatedly over time and observed within a hierarchical trend. The generalized direct blend trend can be effectively adjusted to circumstances where different results are observed. The generalized straight-blend show expects that each result and the univariate models are combined through the detail of dispersion for all arbitrary impacts.

Besides, the blended show can be connected with the detail of negligible dispersion, which is conditional on connected arbitrary impact. Most extreme probabilities and variations are standard for both straight blended models and generalized direct models. This study summarized the effect of fertilizers by utilizing graphic insights such as cruel and standard deviation for ceaseless factors. The generalized straight blend models and model selection criteria were utilized to choose the best-fit models. Akai Information Criteria (AIC) and Bayesian Information Criteria (BIC) were used for model comparison at a 5% level of significance (Hox, Moerbeek and Van de Schoot, 2017; Pongsapakdee and Sukgumphaphan, 2007). Finally, the indicator and reaction factors were imperative for checking the fittingness of the interface (Hosmer Jr., Lemeshow and Sturdivant, 2013).

Model Diagnosis Checking

Residual plots and the Kolmogorov-Simonov test were used for model diagnosis. The residual plots are the best tool for examining whether the selected model meets the normality assumption of the error term or not. A conditional residual plot is important to know the difference between the conditional fits and the observed values in the sample data. Additionally, Kolmogorov-Smirnov test was used to check goodness of fit (Massey Jr, 1951; Wang *et al.*, 2018).

Results

As illustrated in table 1, the mean and standard deviation of the marketable yield of potatoes at the first, second, and third replications of no fertilizer used are (11, SD = 6), (17, SD = 9), and (9, SD = 3) tons per hectare, respectively. In addition, the mean and standard deviation of the marketable yield of potatoes at the first, second, and third replications of NP fertilizer used are (25, SD = 10), (19, SD = 4), and (16, SD = 11) tons per hectare, respectively. Furthermore, the mean and standard deviation of the marketable yield of potatoes at the first, second, and third replications of the NP, S, Zn and B fertilizer used were (25, SD = 6), (23, SD = 7), and (21, SD = 3) tons per hectare, respectively. Therefore, it is possible to conclude that the most important fertilizers obtaining a higher yield of potatoes were N, P, S, Zn, B and NP.

Table 1: Summary Statistics for Marketable Yield within Replication

Independent Variables		Marketable yields (ton/ha)					
		Replication of the experiments					
		First		Second		Third	
		Mean	SD	Mean	SD	Mean	SD
Treatments	No Fertilizer	11	6	17	9	9	3
	NP 165, 195	25	10	19	4	16	11
	NPS 120, 92, 17	19	3	19	9	23	3
	NPSZnB 120, 92, 20.06, 5.89, 0.66	25	6	23	7	21	3
	NPKSZnB 120, 92, 52.17, 19.74, 6.06, 0.88	24	7	20	0	21	4

Table 2 indicates the mean for plant stature, plants per plot, plants per slope, tubers per hill, and marketable yields, which are 73.65 (SD = 12.2), 155.2 (SD = 112.2), 3.23 (SD = 1.31), and 10.91 (SD = 3.15.2), respectively. However, the overall mean of the marketable yield of potatoes is 19.47 (SD = 7.18) tons per hectare. A line graph visualized the relationship between the marketable yield of potatoes and fertilizer types within each replication. It shows the pattern of potato yield measurements over time and the overall individual plots. It is evident from Figure 1 that the variation within and between plots increased slightly with each replication across the plots. The yield of potatoes is slightly turned down when the farmers use NP, NPS, and NPSZnB throughout the first, second, and third replications, respectively.

Table 2: Summary statistics for covariates included in this Study

<i>Covariates</i>	<i>N</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Mean</i>	<i>Std. Deviation</i>
Plant height (cm)	43	51	99	73.65	12.200
Number of plants per plot (count)	43	12	383	155.23	112.227
Number of plants per hill (count)	43	1	6	3.23	1.306
Number of tubers per hill (count)	43	4	17	10.91	3.146
Marketable yields (ton/ha)	43	6	36	19.47	7.182
Valid N (list wise)	43				




Figure 1: Line graph of marketable yield of potato for each fertilizer

The residual diagnostics of the fitted model for the distinction between standardization and scaling of residuals is shown in figure 2. The conditional residual plots require iterative influence analysis or a profiled residual variance and the residuals and the fitted values confirm linearity without distinct patterns (see Figure 2a, and 2d), i.e., constant variance. The quantile-normal plot approves the normality of errors; residual points follow the straight dashed line (Figure 2b and 2c). On the other hand, since the study used the Kolmogorov-Smirnov test for goodness of fit tests, it shows normality with a p-value equal to 0.150.

Table 3 indicates that the random intercept model is well fitted to the data, which has a small AIC, BIC and generalized Chi-square/df values. Hence, this study used a random intercept model to estimate parameters of the associated factors towards potato yields in Basketo Special Woreda, Gamogofa Zone, Ethiopia. The table 4 portrays the intra-class relationship, and it was found to be $(1.4792/1.4792+23.3129) = 0.05992$. 6% of the variation in the marketable yield of potatoes existed between plots, and the remaining 94% inside plots.

In this manner, it appears that the attractive abdicate of potato generation is decided by fertilizer combination or types, the number of plants per plot, and the number of tubers per slope covariates. From table 5, considering fertilizer types as a predictor variable, compared to no fertilizer used for the production of potatoes, the expected marketable yield of potatoes has a difference between NP fertilizer used and no fertilizer used, as it was about 10.3470 tons per hectare; the difference between NPS fertilizer used and not used was 10.9949 tons per hectare. In addition, the difference between NPSZnB fertilizer used and not used

was 13.5530 tons per hectare. Moreover, the variance between NPKSZnB fertilizer used and not used was 13.0592 tons per hectare per replication, as the result indicates.




Figure 2: Conditional Residual Plots for Model Diagnosis Checking

Table 3: Model Selection

<i>Random effects</i>	<i>AIC</i>	<i>BIC</i>	<i>Generalized Chi-Square / df</i>
Intercept only model	296.5	293.7	50.2
Random intercept model	281.4	271.5	23.31
Random coefficients model	291.5	286.1	38.9

Table 4: Estimation of Covariance Parameter for the Marketable Yield of Potato

<i>Covariance Parameters</i>	<i>Subject</i>	<i>Estimate</i>	<i>Standard Error</i>
Variance	Replication	1.4792	2.7055
AR (1)	Replication	0	
Residual		23.3129	5.2341

In other words, the marketable yield of potato changes for each NP fertilizer used was 10.347 times that of the non-fertilizer used in each replication. When NPS fertilizer was used in each replication, the marketable yield of potatoes was 10.9949 times greater than when no fertilizer was used. Additionally, the marketable yield of potatoes for NPSZnB in each replication was 13.55 times that of the amount of fertilizer not used. In expansion, the rate of change of the marketable yield of potatoes for NPKSZnB in each replication was 13.0592 times higher than without including fertilizer used to keep the other factors consistent. This clearly shows that the expected marketable yield of potatoes is increased by 0.0175 tons per hectare when the number of plants per plot is increased by 1 unit. However, the potato yields are decreased by 0.7819 tons/hectare when the number of tubers per slope was increased by 1 unit.

Table 5: Parameter Estimation of Random Intercept Model

<i>Variables</i>	<i>levels</i>	<i>Estimate</i>	<i>SE</i>	<i>P-value</i>	<i>95% CI</i>	
Intercept	Constant	19.1100	5.7302	0.0794	-5.5449	43.7649
Treatments	NPKSZnB	13.0592	2.9009	<.0001	7.1572	18.9612
	NP	10.3470	2.6864	0.0005	4.8814	15.8126
	NPS	10.9949	2.7339	0.0003	5.4328	16.5570
	NPSZnB	13.5530	2.7840	<.0001	7.8890	19.2170
	Not used	0
Plant height (cm)	Continues	-0.04529	0.09712	0.6440	-0.2429	0.1523
Number of plants per plot	Count	0.01750	0.007500	0.0259	0.002242	0.03276
Number of tubers per hill	Count	-0.7819	0.3664	0.0404	-1.5274	-0.03646

Discussion

The random intercepts model shows that the attractive abdicate of potato generation is decided by fertilizer sorts, the number of plants per plot, and the number of tubers per slope covariates. Additionally, 6% of the variation in the marketable yield of potatoes existed between plots, and the remaining 94% of the variety existed inside plots. This finding shows that the fertilizer type and rate were significantly associated with the marketable yield of potatoes. The productivity of potatoes in Ethiopia is very low due to improper use of fertilizer. Potatoes are the most important crops for both cash and food security in developing countries like Ethiopia. This result was confirmed by a study done in Ethiopia (Chindi *et al.*, 2017; Oumer, Hjortsø and de Neergaard, 2013).

This study shows the effect of proper use of fertilizer types on reducing the unmarketable yield of potatoes by increasing the marketable yield of potatoes. The marketable yields of potatoes are increased by 10.3470, 10.9949, 13.5530, and 13.0592 tons per hectare when the farmer used NP, NPS, NPSZnB, and NPKSZnB fertilizer types, respectively. These results are in agreement with the study carried out by various researchers (Al-Taey, Al-Naely and Kshash, 2019). Based on the study done in Russia, Zewide, Singh and Kassa (2021) stated that the application of a combination of 75% of mineral NP can significantly increase the tuber yield. The finding that was reported in Egypt showed significant increases in the total and marketable yield of potato crops from plots treated with 50% of the recommended mineral fertilizers (El-Sayed, Hassan and El-Mogy, 2015). These results also coincide with the study done in Egypt by Ali *et al.* (2022) showing that the increase in the total yield of tubers may be due to the increase in the number of tubers, promoting nutrient uptake and enhancing plant growth through its ability to produce plant hormones as a result of blended or combined fertilizers. Generally, the overall finding indicates that the marketable yield of potatoes increased by 0.0175 tons per hectare when the number of plants per plot increases by 1 unit. Similar findings are conveyed by other researchers (Manea, Al-Bayati and Al-Taey, 2019; Shahnazari *et al.*, 2007). In addition,

the marketable yield of potatoes increased by 0.7819 tons/hectare when the number of tubers per hill decreases by 1 unit. Similar results are reported by other scholars (Al-Taey, Al-Naely and Kshash, 2019). The study was done on the relationship between total yield, and the number of tubers in potato crops, and shows the yield of large tubers is linearly related to total yield and the number of tubers; but where tubers are graded by size, the relationship varies with tuber shape. The main mechanisms available to growers for altering the number of tubers is varietal choice and seed rate but these are also likely to alter total yield with an unpredictable net effect (Burstall, Thomas and Allen, 1987).

Conclusion

Enhanced farm productivity is basic to food security and reducing poverty in developing countries, including Ethiopia. The purpose of this study is to evaluate the contribution of varied fertilizers to potato yields at Basketo Special Woreda, Gamogofa Zone, Ethiopia. To address this objective, a generalized linear mixed model was used to differentiate the factors that affect the marketable yield of potatoes by considering variability within, and between plots. The random intercept model is well fitted to this data, and it shows fertilizer types, number of plants per plot, and number of tubers per hill was significant at 5% level of significance. The expected marketable yield of potatoes increased by 0.7819 when the number of tubers per hill increased by 1 unit. Additionally, the expected marketable yield of potatoes increased by 0.0175 tons per hectare when the number of plants per plot increased by 1 unit. The result of this study suggests that the Ethiopian agriculture authorities should encourage farmers to use NP, NPS, NPSZnB, and NPKSZnB fertilizer types. The total yield of tubers increased may be due to the increase in the number of tubers, promoting nutrient uptake and enhancing plant growth through its ability to produce plant hormones or soil quality. Therefore, the variation of the marketable yield of potatoes within plots was higher than between plots due to the quality of the soil and quantity of fertilizers.

Hence, potato productivity of farmers is basic in food security and reduces poverty. Therefore, government and non-governmental organizations as well as other stakeholders should provide awareness to the farmers for fertilizers rate and potato production. The researchers should also give attention to compare NP, NPS, NPSZnB, and NPKSZnB fertilizers with natural fertilizers, seasonal variation and potato varieties.

Acknowledgments

This field experiment and research by Arba Minch Agricultural Research Center and Southern Agricultural Research Institute, as well as all staff members, are gratefully acknowledged. The authors also acknowledge the Scribes Free Grammar Checker Teams for help regarding English grammar checking.

References

- Alemayehu, M. and Jemberie, M. (2018). Optimum rates of NPS fertilizer application for economically profitable production of potato varieties at Koga Irrigation Scheme, Northwestern Ethiopia. *Cogent Food & Agriculture*, 4(1): 1439663. DOI: <https://doi.org/10.1080/23311932.2018.1439663>.
- Ali, A.M., Hegab, S.A., Abd El Gawad, A. M. and Awad, M. (2022). Integrated Effect of Filter Mud Cake Combined with Chemical and Biofertilizers to Enhance Potato Growth and Its Yield. *Journal of Soil Science and Plant Nutrition*, 22(1): 455-464. DOI: <https://doi.org/10.1007/s42729-021-00661-3>.
- Al-Taey, D.K., Al-Naely, I.J. and Kshash, B.H. (2019). A study on effects of water quality, cultivars, organic and chemical fertilizers on potato (*Solanum tuberosum* L.) growth and yield to calculate the economic feasibility. *Bulgarian Journal of Agricultural Science*, 25(6): 1239-1245. DOI: <https://doi.org/10.1088/1755-1315/910/1/012063>.

- Bellele, T. (2015). Soil fertility mapping and fertilizer recommendation in Ethiopia: Update of Ethiopia SIS project and status of fertilizer blending plants. IPI-MoANR-ATA-Hawassa University Joint Symposium. The role of potassium in balanced fertilization (pp. 24-26). Available online at: <https://www.ipipotash.org/uploads/udocs/9-tegbaru-bellele-soil-fertility-mapping-and-fertilizer-recommendation-in-ethiopia.pdf> [Accessed 11 September 2021].
- Briefs, E.A.M. (2019). Organic farming in the EU a fast-growing sector. Available online at: http://ec.europa.eu/agriculture/markets-and-prices/market-briefs/index_en.htm [Accessed 11 September 2021].
- Burstall, L., Thomas, M.N. and Allen, E.J. (1987). The relationship between total yield, number of tubers and yield of large tubers in potato crops. *The Journal of Agricultural Science*, 108(2): 403-406. DOI: <http://dx.doi.org/10.1017/S0021859600079430>.
- Chindi, A., Shunka, E., Solomon, A., Gebremedhin, W.G., Seid, E. and Tessema, L. (2017). Participatory potato seed production: a breakthrough for food security and income generation in the central highlands of Ethiopia. *Open Agriculture*, 2(1): 205-212. DOI: <https://doi.org/10.1515/opag-2017-0021>.
- El-Sayed, S.F., Hassan, H.A. and El-Mogy, M.M. (2015). Impact of bio-and organic fertilizers on potato yield, quality and tuber weight loss after harvest. *Potato Research*, 58(1): 67-81. DOI: <https://doi.org/10.1007/s11540-014-9272-2>.
- Girma, T., Beyene, S. and Biazin, B. (2017). Effect of organic and inorganic fertilizer application on soil phosphorous balance and phosphorous uptake and use efficiency of potato in Arbegona District, Southern Ethiopia. *Journal of Fertilizers & Pesticides*, 8: 1-6. DOI: <https://doi.org/10.4172/2471-2728.1000185>.
- Girma, T., Beyene, S., Lemaga, B. and Temesgen, B.B. (2020). Integrated application of organic and blended mineral fertilizers improves potato productivity and income for smallholder farmers in acidic soils. *Environment and Natural Resources Research*, 10(1): 61-70. DOI: <https://doi.org/10.5539/enrr.v10n1p61>.
- Hawkesford, M.J. and De Kok, L.J. (2006). Managing sulphur metabolism in plants. *Plant, Cell & Environment*, 29(3): 382-395. DOI: <https://doi.org/10.1111/j.1365-3040.2005.01470.x>.
- Hosmer Jr, D.W., Lemeshow, S. and Sturdivant, R.X. (2013). *Applied logistic regression* (Vol. 398). London: John Wiley & Sons. DOI: <https://doi.org/10.1111/j.1365-3040.2005.01470.x>.
- Hox, J.J., Moerbeek, M. and Van de Schoot, R. (2017). *Multilevel analysis: Techniques and applications*. London: Routledge. DOI: <https://doi.org/10.4324/9781315650982>.
- Manea, A.I., Al-Bayati, H.J. M. and Al-Taey, D.K. (2019). Impact of yeast extract, zinc sulphate and organic fertilizers spraying on potato growth and yield. *Research on Crops*, 20(1): 95-100. DOI: <https://doi.org/10.31830/2348-7542.2019.013>.
- Massey Jr, F.J. (1951). The Kolmogorov-Smirnov test for goodness of fit. *Journal of the American Statistical Association*, 46(253): 68-78. DOI: <https://doi.org/10.1080/01621459.1951.10500769>.
- Oumer, A.M., Hjortsø, C.N. and de Neergaard, A. (2013). Understanding the relationship between livelihood strategy and soil management: Empirical insights from the central highlands of Ethiopia. *Food Security*, 5(2): 143-156. DOI: <https://doi.org/10.1007/s12571-013-0242-3>.
- Pongsapakdee, V. and Sukgumphaphan, S. (2007). Goodness of fit of cumulative logit models for ordinal response categories and nominal explanatory variables with two-factor interaction. *Science, Engineering and Health Studies*, 1(2): 29-38. Available online at: <https://li01.tci-thaijo.org/index.php/sehs/article/view/7259> [Accessed 11 November 2021].
- Rens, L.R., Zotarelli, L., Rowland, D.L. and Morgan, K.T. (2018). Optimizing nitrogen fertilizer rates and time of application for potatoes under seepage irrigation. *Field Crops Research*, 215: 49-58. DOI: <https://doi.org/10.1016/j.fcr.2017.10.004>.
- Rodríguez, F., Ghislain, M., Clausen, A.M., Jansky, S.H. and Spooner, D.M. (2010). Hybrid origins of cultivated potatoes. *Theoretical and Applied Genetics*, 121(6): 1187-1198. DOI: <https://doi.org/10.1007/s00122-010-1422-6>.

- Rosen, C.J., Kelling, K.A., Stark, J.C. and Porter, G.A. (2014). Optimizing phosphorus fertilizer management in potato production. *American Journal of Potato Research*, 91(2): 145-160. DOI: <https://doi.org/10.1007/s12230-014-9371-2>
- Sebnie, W., Esubalew, T. and Mengesha, M. (2021). Response of potato (*Solanum tuberosum* L.) to nitrogen and phosphorus fertilizers at Sekota and Lasta districts of Eastern Amhara, Ethiopia. *Environmental Systems Research*, 10(1): 1-8. DOI: <https://doi.org/10.1186/s40068-020-00213-1>.
- Setu, H., Dechassa, N. and Alemayehu, Y. (2018). Influence of phosphorus and potassium fertilizers on growth and yield of potato (*Solanum tuberosum* L.) at Assosa, Benishangul Gumuz Regional State, Western Ethiopia. *Journal of Soil Science and Environmental Management*, 9(6): 81-90. DOI: <https://doi.org/10.5897/JSSEM2016.0568>.
- Shahnazari, A., Liu, F., Andersen, M.N., Jacobsen, S.E. and Jensen, C.R. (2007). Effects of partial root-zone drying on yield, tuber size and water use efficiency in potato under field conditions. *Field Crops Research*, 100(1): 117-124. DOI: <https://doi.org/10.1016/j.fcr.2006.05.010>.
- Sonnenwald, S., van Harsselaar, J., Ott, K., Lorenz, J. and Sonnewald, U. (2015). How potato plants take the heat? *Procedia Environmental Sciences*, 29: 97. DOI: <https://doi.org/10.1515/opag-2020-0006>.
- Tessema, L., Mohammed, W. and Abebe, T. (2020). Evaluation of potato (*Solanum tuberosum* L.) varieties for yield and some agronomic traits. *Open Agriculture*, 5(1): 63-74. DOI: <https://doi.org/10.1515/opag-2020-0006>.
- Voora, V., Larrea, C. and Bermudez, S. (2020). Global market report: bananas. *International Institute for Sustainable Development*. Available online at: <https://www.fruchtportal.de/media/files/Pdf%20diversen%202020/iisd%20sustainability%20report%20ssi%20global%20market%20report%20banana%20mei%202020.pdf> [Accessed 11 September 2021].
- Wang, Y., Chen, Q., Zhang, N. and Wang, Y. (2018). Conditional residual modeling for probabilistic load forecasting. *IEEE Transactions on Power Systems*, 33(6): 7327-7330. DOI: <https://doi.org/10.1109/TPWRS.2018.2868167>.
- Zewide, I., Singh, S. and Kassa, H. (2021). Potato Productivity Enhancement by Combined Use of Blended Fertilizer (NPSB), Cattle Manure, Vermicompost, and Mineral NP in Acidic Soil. *Russian Agricultural Sciences*, 47(1): S27-S33. DOI: <https://doi.org/10.3103/S1068367422010037>.

Authors' Declarations and Essential Ethical Compliances

Authors' Contributions (in accordance with ICMJE criteria for authorship)

Contribution	Author 1	Author 2	Author 3	Author 4	Author 5
Conceived and designed the research or analysis	Yes	No	Yes	No	No
Data collection	Yes	Yes	No	No	Yes
Contributed to data analysis & interpretation	No	No	Yes	No	No
Written the article/paper	Yes	No	Yes	Yes	No
Critical revision of the article/paper	Yes	No	Yes	Yes	No
Editing of the article/paper	Yes	No	Yes	Yes	No
Supervision	Yes	No	Yes	Yes	No
Project Administration	No	No	Yes	No	No
Funding Acquisition	No	No	No	No	No
Overall Contribution Proportion (%)	25	5	50	15	5

Funding

Haramaya University contributed for the research conducted for writing this paper.

Research involving human bodies (Helsinki Declaration)

Has this research used human subjects for experimentation? No

Research involving animals (ARRIVE Checklist)

Has this research involved animal subjects for experimentation? No

Research involving Plants

During the research, the authors followed the principles of the Convention on Biological Diversity and the Convention on the Trade in Endangered Species of Wild Fauna and Flora. Yes

Research on Indigenous Peoples and/or Traditional Knowledge

Has this research involved Indigenous Peoples as participants or respondents? No

(Optional) PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses)

Have authors complied with PRISMA standards? Yes

Competing Interests/Conflict of Interest

Authors have no competing financial, professional, or personal interests from other parties or in publishing this manuscript.

Rights and Permissions

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.